

## Mini Review

# Doping Behaviors and Prevention in Amateur Sport: An Update and New Perspective

Lentillon-Kaestner V\*

*Teaching and Research Unit in Physical Education and Sport (UER-EPS), University of Teacher Education, State of Vaud (HEP-VD), Switzerland*

## \*Corresponding author

Vanessa Lentillon-Kaestner, Teaching and Research Unit in Physical Education and Sport, University of Teacher Education, State of Vaud, Avenue de Cour 25, CH1014 Lausanne, Switzerland, Tel: 41-21-316-36-16; Email: Vanessa.lentillon-kaestner@hepl.ch

Submitted: 23 March 2015

Accepted: 30 April 2015

Published: 04 May 2015

## Copyright

© 2015 Lentillon-Kaestner

## OPEN ACCESS

## Keywords

- Doping
- Prevention
- Sport
- Fear-based campaigns

## Abstract

**Background:** Doping is not limited to elite athletes. To reduce doping in sport, it seems important to influence young athletes in primary prevention.

**Purpose:** Based on previous research, the purpose of this paper is to give an overview on doping behaviors in amateur sport, actual prevention campaigns, and to propose a new perspective in doping prevention.

**Methods:** This mini-review is based on articles published in English and French between 1990 and 2014 retrieved from different databases (i.e., PubMed, Science Direct, Francis, Cairn. Info). In addition, some edited books and chapter books, and recent research reports were used to complete this mini-review.

**Results:** Doping is increasingly important among amateur athletes. To date, traditional doping prevention campaigns are often ineffective. In recent years, a new model of prevention campaigns based on fear, coming from the Anglo-Saxon and Scandinavian countries, has been used notably in France (e.g., prevention campaigns for road safety, tobacco, alcohol, cancer). Fear-based campaigns have scientific support and have shown a relatively small but still solid effect on attitudes, intentions and behaviors.

**Conclusion:** The fight against doping would benefit from trying fear-based anti-doping campaigns.

## ABBREVIATIONS

EPPM: Extended Parallel Process Model

## INTRODUCTION

Based on previous research, the purpose of this paper is to give an overview of doping behaviors in amateur sport, actual prevention campaigns, and to propose a new perspective in doping prevention. This mini-review is based on articles published in English and French between 1990 and 2014 retrieved from different databases (i.e., PubMed, Science Direct, Francis, Cairn. Info). The following words and combinations of words in English and French have been used on databases: #1 doping, #2 drugs, #3 anabolic steroids, #4 amateur sport, #5 recreational sport, #6 prevention, #7 campaigns, #8 fear, #9 threat, #10 EPPM, #11 self-affirmation, (#1 OR #2 OR #3) AND (#4 OR #5 OR #6 OR #7), (#6 OR #7) and (#8 OR #9 OR #10), (#6 OR #7 OR #8 OR #9 OR #10) AND (#11). In addition, some edited books and chapter books, and recent research reports were used to complete this mini-review. The more significant studies were subjected to narrative synthesis, summarizing the current state

of knowledge in relation to doping behaviors and prevention in amateur sport.

## Widespread doping behaviors among amateur athletes

Doping is not limited to elite athletes but is widespread in society and is increasingly important among amateur athletes [1-20]. It is difficult to assess the extent of doping in amateur sport, nevertheless it exists [17]. In his review on 44 studies, Laure (1997) estimated the prevalence of doping in children and adolescents participating in sport at 3 to 5 % and in adults participating in amateur sports at 5 to 15 % [7]. In France, 6.7 % of 8-18 year-olds approved doping in sport [21]. Young athletes associated positively sport with drug consumption, at least with energizing drugs [22]. In the study of Turblin et al. (1995), 9 % of students practicing sports outside physical education reported having consumed a substance that they considered as doping [4]. Lentillon-Kaestner and Carstairs (2010) showed that young cyclists (Under-23 category) were tempted by doping [16]. The meta-analysis of Sagoe, Molde and Andreassen (2014) on 187 studies showed a global lifetime prevalence rate of anabolic-androgenic steroid use of 3.3 %.

Doping varies according to various demographic parameters. It increases with age and can start before the age of 15 years [3,4,6,7,19,23-25]. Doping is more widespread among boys than girls [4,21,23-26]; however, the gender gap is decreasing from 10 years old [8]. Doping is more widespread among competitors, and it increases with the level of competition [2,4,6,21,25].

### Inefficiency of current anti-doping campaigns

For several years, the fight against doping has mainly focused on the improvement of detection measures (drug tests), leaving aside measures of doping prevention [27-29]. To date, tested anti-doping campaigns are rare, and doping prevention programs lack solid scientific background [27,28,30,31]. Traditional doping prevention campaigns are often ineffective. They describe substances' side effects, try to persuade users of the ineffectiveness of performance enhancing substances or promote sports ethics [28,32]. The recent meta-analysis of Ntoumakis, Ng, Barkoukis and Backhouse (2014) showed that implemented anti-doping interventions lead to small changes in individuals' attitudes towards and intention to engage in doping and had no effect on actual doping behaviors [28]. It seems important to build innovative prevention campaigns that are based on solid scientific theory [9,27,30].

### The fear-based campaigns

In recent years, a new model of prevention campaign based on fear and coming from the Anglo-Saxon and Scandinavian countries has been used notably in France (e.g., road safety, tobacco, alcohol, cancer prevention campaigns) [33]. Fear is conceptualized as a negative emotional reaction to a perceived threat. The purpose of the fear model is to show the consequences of an undesirable event (e.g., illness, accident) or to give more or less directly a glimpse of the following unhappiness aiming to bring an attitude change. The fear motivates actions to reduce negative emotion [34]. In contrast to current doping prevention campaigns, fear-based campaigns have scientific support [35-45]. Psychologists and researchers in marketing have tried to understand why a prevention campaign based on phobic emotion resulted sometimes in success (action) and sometimes in failure (defensive reactions). Various theories have been developed. The latest and most advanced theory about fear appeals from a theoretical and empirical point of view [33,40] is the Extended Parallel Process Model (EPPM) of Witte (1992, 1998) [36,46]. In this model, individuals first assess the threat contained in the message. Perceived threat is a cognitive construct with two dimensions: perceived severity of the threat and one's perceived susceptibility to the threat [47]. In accordance with other meta-analyses, the meta-analysis of Witte and Allen (2000) suggested that the higher the fear level, the higher is the persuasive impact of the message [40]. If the threat is perceived as irrelevant or insignificant, the person is no longer motivated to process the message and simply ignores the fear. In contrast, when a threat is described as significant and relevant, people are frightened. The more people believe themselves vulnerable to a serious threat, the more they are motivated to start the second evaluation of the recommendations' effectiveness. The fear motivates the change in attitudes, intentions and behaviors, especially fear accompanied with highly effective messages. Perceived effectiveness comprises two dimensions: perceived response effectiveness (beliefs of

how effective a response is in averting a threat) and perceived self-effectiveness (beliefs about one's ability to carry out the recommended response) [47]. Effective messages generating a strong fear encourage behavior change (i.e., danger control), while less effective messages generating a strong fear lead to defensive reactions (i.e., fear control) [40,47,48]. According to Witte (1992), fear-based health campaigns are far more useful to promote prevention behaviors than to modify existing behaviors [46]. Witte and Allen (2000) concluded, from their meta-analysis on 98 studies on fear-based campaigns (e.g., sexuality, alcohol, road safety, tobacco), that fear would have a relatively small but constant effect on attitudes, intentions and behaviors [40]. They also offered a series of recommendations for the implementation of prevention measures [40]. In addition, psychology studies on persuasion showed that a simple message was more persuasive in video than in written or audio forms [49,50].

The theory of self-affirmation appears as a way to increase the effectiveness of prevention campaigns through a re-evaluation of the self-image, which reduces the defensive reactions and increases the acceptance of prevention recommendations [51]. The manipulation of self-affirmation may be achieved in different ways (e.g., values to rank in importance order, to write an essay on their most important value, to describe a very important thing in their lives) [52]. Research has shown that to secure the self through self-affirmation manipulation reduced defensive reactions to threatening health information [51] and positively influenced healthier behaviors [53,54]. Through the self-affirmation process, prevention campaigns do not threaten the self-image of the person, but only the behavior is threatened [51,52].

### New perspectives in doping prevention

Adolescence is a high-risk period for the development of doping behaviors [55]. Even if performance enhancing drugs have adverse effects on health [12,29,56-58], young athletes are tempted by doping and are not afraid of health side effects [59,60]. Young athletes are priority target as their doping attitudes are in formation and primary prevention seems to be a good solution to avoid the appearance of doping behaviors.

To date, no previous study has evaluated the effects of fear-based anti-doping campaigns and the fight against doping would benefit from trying a fear-based doping prevention video for two main reasons. Firstly, although very significant progress has been made in detection measures in recent years in the fight against doping, prevention campaigns remain limited. Actual implemented doping prevention measures are ineffective and without scientific support. Secondly, doping prevention lacks standardized, effective and easy tools to use in both the sport and the academic domains. A doping prevention video could be used during sport events and competitions. Doping prevention is also a topic addressed in sport courses (i.e., additional sport option, sports study) in high schools and in universities. A fear-based anti-doping video would be directly useful for all stakeholders (e.g., teachers, coaches, sport and doping prevention organizations) who wish to address this issue with athletes and students. It will prevent the development of prevention campaigns that may have opposite effects than expected. To indicate the main effects of doping substances (which is common in prevention campaigns)

has counterproductive effects on athletes by even increasing their desire to test these substances. Doping is not an easy issue to address. The fear-based doping prevention video would encourage a debate among young people and should reduce the athletes' intentions to use doping.

## REFERENCES

1. Anderson WA, Albrecht RR, McKeag DB, Hough DO, McGrew CA. A national survey of alcohol and drugs use by college athletes. *Physician and Sportsmedicine*. 1991; 19: 91-104.
2. Skowno J. Drug survey among first-team schoolboy rugby players. *S Afr Med J*. 1992; 82: 204.
3. Radakovich J, Broderick P, Pickell G. Rate of anabolic-androgenic steroid use among students in junior high school. *J Am Board Fam Pract*. 1993; 6: 341-345.
4. Turblin P, Grosclaude P, Navarro F, Rivière D, Garrigues M. Enquête épidémiologique sur le dopage en milieu scolaire dans la région Midi-Pyrénées. *Science & Sports*. 1995;10: 87-94.
5. MacAuley D. Drugs in sport. *BMJ*. 1996; 313: 211-215.
6. Melia P, Pipe A, Greenberg L. The use of anabolic-androgenic steroids by Canadian students. *Clin J Sport Med*. 1996; 6: 9-14.
7. Laure P. Epidemiologic approach of doping in sport. A review. *J Sports Med Phys Fitness*. 1997; 37: 218-224.
8. Yesalis CE, Barsukiewicz CK, Kopstein AN, Bahrke MS. Trends in anabolic-androgenic steroid use among adolescents. *Arch Pediatr Adolesc Med*. 1997; 151: 1197-1206.
9. Yesalis CE, Bahrke MS. Doping among adolescent athletes. *Baillieres Best Pract Res Clin Endocrinol Metab*. 2000; 14: 25-35.
10. Pillard F, Grosclaude P, Navarro F, Godeau E, Rivière D. Pratique sportive et conduite dopante d'un échantillon représentatif des élèves de Midi-Pyrénées. *Science & Sports* 2002; 17: 8-16.
11. Laure P, Binsinger C. Adolescent athletes and the demand and supply of drugs to improve their performance. *J Sports Sci Med*. 2005; 4: 272-277.
12. Calfee R, Fadale P. Popular ergogenic drugs and supplements in young athletes. *Pediatrics*. 2006; 117: e577-589.
13. Laure P, Binsinger C. Doping prevalence among preadolescent athletes: a 4-year follow-up. *Br J Sports Med*. 2007; 41: 660-663.
14. Nandrino JL, Escande J-D, Faure S, Doba K, Vandeweege E. Profil psychologique et comportemental de vulnérabilité à la dépendance à l'exercice et au risque de pratiques dopantes chez les sportifs amateurs: l'exemple des semi-marathoniens. *Annales Médico-Psychologiques*. 2008; 166: 772-778.
15. Sjöqvist F, Garle M, Rane A. Use of doping agents, particularly anabolic steroids, in sports and society. *Lancet*. 2008; 371: 1872-1882.
16. Lentillon-Kaestner V, Carstairs C. Doping use among young elite cyclists: a qualitative psychosociological approach. *Scand J Med Sci Sports*. 2010; 20: 336-345.
17. Lentillon-Kaestner V, Ohl F. Can we measure accurately the prevalence of doping? *Scand J Med Sci Sports*. 2011; 21: e132-142.
18. Zelli A, Mallia L, Lucidi F. The contribution of interpersonal appraisals to a social-cognitive analysis of adolescents' doping use. *Psychology of Sport and Exercise*. 2010; 11: 304-311.
19. Sagoe D, Molde H, Andreassen CS, Torsheim T, Pallesen S. The global epidemiology of anabolic-androgenic steroid use: a meta-analysis and meta-regression analysis. *Ann Epidemiol*. 2014; 24: 383-398.
20. Barkoukis V, Lazuras L, Harris PR. The effects of self affirmation manipulation on decision making about doping use in elite athletes. *Psychology of Sport and Exercise*. 2015; 16: 1-7.
21. Laure P. *Dopage et société*. Paris: Ellipses; 2000.
22. CNRS. *Dopage et pratiques sportives*. Paris: Expertise collective CNRS; 1998.
23. Dunn MO, Thomas J. A risk profile of elite Australian athletes who use illicit drugs. *Addict Behav*. 2012; 37: 144-147.
24. Scott DM, Wagner JC, Barlow TW. Anabolic steroid use among adolescents in Nebraska schools. *Am J Health Syst Pharm*. 1996; 53: 2068-2072.
25. Pillard F, Grosclaude P, Navarro F, Godeau E, Rivière D. Enquête épidémiologique sur le dopage sportif en milieu scolaire dans la région midi-pyrénées en 1999. Résultats préliminaires. *Bulletin Epidémiologique Hebdomadaire (BEH)*. 2000; 42: 185-186.
26. Mignon P. *Le dopage: état des lieux sociologiques*. Paris: Université Paris 5 René Descartes, 2002.
27. Backhouse SH, Patterson, L, McKenna, J. Achieving the Olympic ideal: preventing doping in sport. *Performance Enhancement & Health*. 2012; 1: 83-85.
28. Ntoumanis N, Ng JY, Barkoukis V, Backhouse S. Personal and psychosocial predictors of doping use in physical activity settings: a meta-analysis. *Sports Med*. 2014; 44: 1603-1624.
29. Overbye M, Knudsen, M.L, Pfister, G. To dope or not to dope: Elite athletes' perceptions of doping deterrents and incentives. *Performance Enhancement & Health*. 2013; 2: 119-134.
30. Johnson MB. A systemic social cognitive perspective on doping. *Psychology of sport and exercise*. 2012; 13: 317-323.
31. Moston S, Engelberg, T, Skinner, J. Self fulfilling prophecy and the future of doping. *Psychology of Sport and Exercise*. 2015; 16: 201-207.
32. Laure P, Lecerf T. Prévention du dopage chez les adolescents: comparaison d'une démarche éducative et d'une démarche informative. *Science & Sports*. 2002; 17: 198-201.
33. Gallopel K, Petr C. Utilisation de la peur dans les campagnes de prévention : résultats et discussions autour des comportements tabagiques de jeunes français. *Convegno Le Tendenze del marketing in Europa; Venise; 2000*.
34. Gallopel K. L'utilisation de la peur dans un contexte de marketing social: état de l'art, limites et voies de recherche. *Recherche et Application en Marketing*. 2006; 21: 41-60.
35. Witte K. Fear control and danger control: a test of the extended parallel process model (EPPM). *Communication Monographs*. 1994; 61: 113-134.
36. Witte K, Berkowitz JM, Cameron KA, McKeon JK. Preventing the spread of genital warts: using fear appeals to promote self-protective behaviors. *Health Educ Behav*. 1998; 25: 571-585.
37. Moscato S, Black DR, Blue CL, Mattson M, Galer-Uni RA, Coster DC. Evaluating a fear appeal message to reduce alcohol use among "Greeks". *Am J Health Behav*. 2001; 25: 481-491.
38. Tay R, Watson B. Changing drivers intentions and behaviours using fear-based driver fatigue advertisements. *Health Mark Q*. 2002; 19: 55-68.
39. Smalec JL, Klingle RS. Bulimia interventions via interpersonal influence: the role of threat and efficacy in persuading bulimics to seek help. *J Behav Med*. 2000; 23: 37-57.
40. Witte K, Allen M. A meta-analysis of fear appeals: implications for

- effective public health campaigns. *Health Educ Behav.* 2000; 27: 591-615.
41. McKay DL, Berkowitz JM, Blumberg JB, Goldberg JP. Communicating cardiovascular disease risk due to elevated homocysteine levels: Using the EPPM to develop print materials. *Health Education & Behavior.* 2004; 31: 355-371.
42. Smith SW, Rosenman KD, Kotowski MR, Glazer E, McFeters C, Keesecker NM, et al. Using the EPPM to create and evaluate the effectiveness of brochures to increase the use of hearing protection in farmers and landscape workers. *Journal of Applied Communication Research.* 2008; 36: 200-218.
43. Wong NC, Cappella JN. Antismoking Threat and Efficacy Appeals: Effects on Smoking Cessation Intentions for Smokers with Low and High Readiness to Quit. *J Appl Commun Res.* 2009; 37: 1-20.
44. Cho H, Witte K. Managing fear in public health campaigns: a theory-based formative evaluation process. *Health Promot Pract.* 2005; 6: 482-490.
45. Morrison K. Motivating women and men to take protective action against rape: examining direct and indirect persuasive fear appeals. *Health Commun.* 2005; 18: 237-256.
46. Witte K. Putting the fear back into fear appeals: the Extended Parallel Process Model. *Communication Monographs.* 1992; 59: 329-349.
47. Popova L. The extended parallel process model: illuminating the gaps in research. *Health Educ Behav.* 2012; 39: 455-473.
48. Ruitter R, Verplanken B, De Cremer D, Kok G. Danger and fear control in response to fear appeals: the role of need for cognition. *Basic and Applied Social Psychology.* 2004; 26: 13-24.
49. Girandola F. *Peur et persuasion: présentations des recherches (1953-1998) et d'une nouvelle lecture. L'Année Psychologique.* 2000; 100: 333-376.
50. Girandola F. *Psychologie de la persuasion et de l'engagement: Presses Universitaires de Franche-Comté; 2003.*
51. Sherman DAK, Nelson LD, Steele CM. Do messages about health risks threaten the self? Increasing the acceptance of threatening health messages via self-affirmation. *Personality and Social Psychology Bulletin.* 2000; 26: 1046-1058.
52. McQueen A, Klein WMP. Experimental manipulations of self-affirmation: a systematic review. *Self-identity.* 2006; 5: 289-354.
53. Harris PR. Self-affirmation and the self-regulation of health behavior change. *Self and Identity.* 2011; 10: 304-314.
54. Harris PR, Epton T. The impact of self-affirmation on health cognition, health behavior and other health-related responses: a narrative review. *Social and Personality Compass.* 2009; 3: 962-978.
55. Schirlin O, Rey G, Jouvent R, Dubal S, Komano O, Perez-Diaz F, et al. Attentional bias for doping words and its relation with physical self-esteem in young adolescents. *Psychology of Sport and Exercise.* 2009; 10: 615-620.
56. Maravelias C, Dona A, Stefanidou M, Spiliopoulou C. Adverse effects of anabolic steroids in athletes. A constant threat. *Toxicol Lett.* 2005; 158: 167-175.
57. Grossi O, G n reau T. Glucocorticoids and infections, doping, surgery, sexuality. *Rev Med Interne.* 2013; 34: 269-278.
58. Dine G, Van Lierde, F, Rehn, Y, Gillier, N. Probl mes associ s   l'usage de l' rythropo tine en milieu sportif. *Revue G n rales et Analyses Prospectives.* 2001: 235-243.
59. Irving LM, Wall M, Neumark-Sztainer D, Story M. Steroid use among adolescents: findings from Project EAT. *J Adolesc Health.* 2002; 30: 243-252.
60. Lentillon-Kaestner V, Hagger MS, Hardcastle S. Health and doping in elite-level cycling. *Scand J Med Sci Sports.* 2012; 22: 596-606.

**Cite this article**

Lentillon-Kaestner V (2015) Doping Behaviors and Prevention in Amateur Sport: An Update and New Perspective. A Pilot Study. *Ann Sports Med Res* 2(4): 1029.